



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

River. The Farallones (Spanish, meaning pointed rocks in the sea) are disposed in three groups several miles apart, the largest being about a mile long and lying thirty miles west of the Golden Gate. They are well named, for there is neither soil nor vegetation upon them, except the guano of the birds and three species of weeds. In summer the eggs of the birds which swarm there to breed, are gathered by the barrel-full for the San Francisco market.—CHAS. H. TOWNSEND, *Smithsonian Institution, Washington, D. C.*

The Yellow-rumped Warbler Wintering in Maine.—On January 1, 1885, I shot two Yellow-rumped Warblers (*Dendraeca coronata*) from a flock of six at Pine Point, Maine. On opening the crop of one, I found it filled with the seeds of the pitch pine. I believe this species has never before been taken in the winter season north of Massachusetts.—JOSEPH L. GOODALE, *Cambridge, Mass.*

The Migration of the Swallows.—I have noticed for several years that before the young Swallows were capable of enduring a prolonged flight, old and young gathered together in one vast assembly and moved gradually southward, making short stages from farm to farm; at last (in 1884, on August 9), with a favorable north wind and a clear sky, they left the Island in a body, only a few stragglers remaining, just enough to remind us that summer was still with us.—FRANCIS BAIN, *North River, P. E. I.*

Nelson's Sharp-tailed Finch (*Ammodramus caudacutus nelsoni*) on the Atlantic Coast.—Mr. Arthur T. Wayne sends me a Sharp-tailed Finch which is positively indistinguishable from Illinois specimens, but which was shot on the salt marshes near Charleston, South Carolina, Oct. 8, 1884. That it is really an inland-bred bird scarcely admits of a doubt, nor is its occurrence on this coast altogether surprising in view of the fact that other species which breed only in the interior—*Coturniculus lecontei*, for example—extend their autumnal migrations in a south-easterly direction and winter numerous very near to, if not actually on, the Atlantic seaboard.—WILLIAM BREWSTER, *Cambridge, Mass.*

Wintering of the Swamp Sparrow in Eastern Massachusetts.—The capture of two Swamp Sparrows (*Melospiza palustris*) in Cambridge, on January 11, 1883, has already been recorded,* but a second instance may be of interest.

On December 29, 1884, a flock of four were seen and one killed in a dense thicket on the Fresh Pond marshes in Cambridge, and on January 31, 1885, near the same place, I saw the remains of another, which had been partly eaten by a Shrike. Since then I have looked for them several times unsuccessfully, but think that the rest had probably been killed by Shrikes.—ARTHUR P. CHADBOURNE, *Cambridge, Mass.*

* Journal Boston Zoölogical Society, Vol. II (1883), p. 32.